

DARK ENERGY AND DARK MATTER AS CURVATURE EFFECTS

S. CAPOZZIELLO

*Dipartimento di Fisica, Università di Napoli "Federico II" and
INFN, Sez. di Napoli, Napoli, Italy
E-mail: capozziello@na.infn.it*

Abstract. Extended Theories of Gravity, in particular $f(R)$ gravity, could be, in principle, able to explain the accelerated expansion of the Universe without adding unknown forms of dark energy/dark matter but, more simply, extending the General Relativity by generic functions of the Ricci scalar. However, apart from several phenomenological models, there is no final extended theory capable of fitting all the observations and addressing all the issues related to the presence of dark energy and dark matter. Astrophysical observations are pointing out huge amounts of "dark matter" and "dark energy" needed to explain the observed large scale structures and cosmic accelerating expansion. Up to now, no experimental evidence has been found, at fundamental level, to explain such mysterious components. The problem could be completely reversed considering dark matter and dark energy as "shortcomings" of General Relativity.

1. INTRODUCTION

Although being the best fit to a wide range of data, the Λ CDM model is affected by strong theoretical shortcomings that have motivated the search for alternative models (Copeland et al. 2006). Dark Energy (DE) models mainly rely on the implicit assumption that Einstein's General Relativity (GR) is the correct theory of gravity. Nevertheless, its validity on the larger astrophysical and cosmological scales has never been tested, and it is therefore conceivable that both cosmic speed up and Dark Matter (DM) represent signals of a breakdown of GR. Following this line of thinking, the choice of a generic function $f(R)$ as the gravitational Lagrangian, where R is the Ricci scalar, can be derived by matching the data and by the "economic" requirement that no exotic ingredients have to be added. This is the underlying philosophy of what is referred to as $f(R)$ gravity (Capozziello & De Laurentis 2011). It is worth noticing that Solar System experiments show the validity of GR at these scales so that $f(R)$ theories should not differ too much from GR at this level. In other words, the PPN limit of such models must not violate the experimental constraints on Eddington parameters. A positive answer to this request has been recently achieved for several $f(R)$ theories (Capozziello & Troisi 2005), nevertheless it has to be remarked that this debate is far to be definitively concluded. Although higher order gravity theories have received much attention in cosmology, since they are naturally able to give rise to the accelerating expansion (both in the late and in the early universe, it is possible

to demonstrate that $f(R)$ theories can also play a major role at astrophysical scales (Capozziello et al. 2007, Capozziello et al. 2009). In fact, modifying the gravity action can affect the gravitational potential in the low energy limit. Provided that the modified potential reduces to the Newtonian one on the Solar System scale, this implication could represent an intriguing opportunity rather than a shortcoming for $f(R)$ theories. In fact, a corrected gravitational potential could offer the possibility to fit galaxy rotation curves without the need of Dark Matter. In addition, one could work out a formal analogy between the corrections to the Newtonian potential and the usually adopted Dark Matter models. In order to investigate the consequences of $f(R)$ theories on both cosmological and astrophysical scales, let us first remind the basics of this approach and then discuss dark energy and dark matter issues as curvature effects.

2. DARK ENERGY AS A CURVATURE EFFECT

From a mathematical viewpoint, $f(R)$ theories generalize the Hilbert-Einstein Lagrangian as $\mathcal{L} = \sqrt{-g}f(R)$ without assuming *a priori* the functional form of Lagrangian density in the Ricci scalar. The field equations are obtained by varying with respect to the metric components to get:

$$f'(R)R_{\alpha\beta} - \frac{1}{2}f(R)g_{\alpha\beta} = f'(R)^{;\mu\nu} (g_{\alpha\mu}g_{\beta\nu} - g_{\alpha\beta}g_{\mu\nu}) + T_{\alpha\beta}^M \quad (1)$$

where the prime denotes derivative with respect to the argument and $T_{\alpha\beta}^M$ is the standard matter stress-energy tensor. Defining the *curvature stress-energy tensor* as

$$T_{\alpha\beta}^{curv} = \frac{1}{f'(R)} \left\{ \frac{1}{6}g_{\alpha\beta} [f(R) - Rf'(R)] + f'(R)^{;\mu\nu} (g_{\alpha\mu}g_{\beta\nu} - g_{\alpha\beta}g_{\mu\nu}) \right\}. \quad (2)$$

Eq.(1) may be recast in the Einstein-like form as:

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R = T_{\alpha\beta}^{curv} + T_{\alpha\beta}^M/f'(R) \quad (3)$$

where matter non-minimally couples to geometry through the term $1/f'(R)$. The presence of term $f'(R)_{;\mu\nu}$ renders the equations of fourth order, while, for $f(R) = R$, the curvature stress-energy tensor $T_{\alpha\beta}^{curv}$ identically vanishes and Eqs.(3) reduce to the standard second-order Einstein field equations. As it is clear, from Eq.(3), the curvature stress-energy tensor formally plays the role of a further source term in the field equations so that its effect is the same as that of an effective fluid of purely geometrical origin. However the metric variation is just one of the approaches towards $f(R)$ gravity: in fact, one can face the problem also considering the so called Palatini approach where the metric and connection fields are considered independent. Apart from some differences in the interpretation, one can deal with a fluid of geometric origin in this case as well. The scheme outlined above provides all the ingredients we need to tackle with the dark side of the Universe. Depending on the scales, such a curvature fluid can play the role of DM and DE. From the cosmological point of view, in the standard framework of a spatially flat homogenous and isotropic Universe, the cosmological dynamics is determined by its energy budget through the Friedmann




Figure 1: The Hubble diagram of 20 radio galaxies together with the “gold” sample of SNeIa, in term of the redshift as suggested in Daly & Diorgovsky 2004. The best fit curve refers to the $f(R)$ -gravity model without Dark Matter.

equations. The cosmic acceleration is achieved when the r.h.s. of the acceleration equation remains positive (in physical units with $8\pi G = c = 1$):

$$\frac{\ddot{a}}{a} = -\frac{1}{6}(\rho_{tot} + 3p_{tot}) , \quad (4)$$

where a is the scale factor, $H = \dot{a}/a$ the Hubble parameter, the dot denotes derivative with respect to cosmic time, and the subscript tot denotes the sum of the curvature fluid and the matter contribution to the energy density and pressure. From the above relation, the acceleration condition, for a dust dominated model, leads to:

$$\rho_{curv} + \rho_M + 3p_{curv} < 0 \rightarrow w_{curv} < -\frac{\rho_{tot}}{3\rho_{curv}} \quad (5)$$

so that a key role is played by the effective quantities:

$$\rho_{curv} = \frac{8}{f'(R)} \left\{ \frac{1}{2} [f(R) - Rf'(R)] - 3H\dot{R}f''(R) \right\} , \quad (6)$$

and

$$w_{curv} = -1 + \frac{\ddot{R}f''(R) + \dot{R} [\dot{R}f'''(R) - Hf''(R)]}{[f(R) - Rf'(R)]/2 - 3H\dot{R}f''(R)} . \quad (7)$$

As a first simple choice, one may neglect ordinary matter and assume a power-law form $f(R) = f_0 R^n$, with n a real number, which represents a straightforward generalization of the Einstein GR in the limit $n = 1$. One can find power-law solutions for $a(t)$ providing a satisfactory fit to the SNeIa data and a good agreement with the estimated age of the Universe in the range $1.366 < n < 1.376$ (Capozziello et. 2003). The data fit turns out to be significant (see Fig. 1) improving the χ^2 value and, it fixes the best fit value at $n = 3.46$ when it is accounted that only the baryon

contribute $\Omega_b \approx 0.04$ (according with BBN prescriptions). It has to be remarked that considering DM does not modify the result of the fit, supporting the assumption of no need for DM in this model. From the evolution of the Hubble parameter in term of redshift one can even calculate the Age of Universe. The best fit value $n = 3.46$ provides $t_{univ} \approx 12.41$ Gyr. It is worth noting that considering $f(R) = f_0 R^n$ gravity represents only the simplest generalization of Einstein theory. In other words, it has to be considered that R^n -gravity represents just a working hypothesis as there is no overconfidence that such a model is the correct final gravity theory. In a sense, we want only to suggest that several cosmological and astrophysical results can be well interpreted in the realm of a power law extended gravity model. This approach gives no rigidity about the value of the power n , although it would be preferable to determine a model capable of working at different scales. Furthermore, we do not expect to be able to reproduce the whole cosmological phenomenology by means of a simple power law model, which has been demonstrated not to be sufficiently versatile. For example, we can demonstrate that this model fails when it is analyzed with respect to its capability of providing the correct evolutionary conditions for the perturbation spectra of matter overdensity δ . This point is typically addressed as one of the most important issues which suggest the need for Dark Matter. In fact, if one wants to discard this component, it is crucial to match the observational results related to the Large Scale Structure of the Universe and the Cosmic Microwave Background which show, respectively at late time and at early time, the signature of the initial matter spectrum. As important remark, we note that the quantum spectrum of primordial perturbations, which provides the seeds of matter perturbations, can be positively recovered in the framework of R^n -gravity. In fact, $f(R) \propto R^2$ can represent a viable model with respect to CMBR data and it is a good candidate for cosmological Inflation. To develop the matter power spectrum suggested by this model, we resort to the equation for the matter contrast obtained in Zhang 2006, in the case of fourth order gravity. This equation can be deduced considering the conformal Newtonian gauge for the perturbed metric:

$$ds^2 = (1 + 2\psi)dt^2 - a^2(1 + 2\phi)\Sigma_{i=1}^3(dx^i). \quad (8)$$

In GR, it is $\phi = -\psi$, since there is no anisotropic stress; in extended gravity, this relation breaks, in general, and the $i \neq j$ components of field equations give new relations between ϕ and ψ . In particular, for $f(R)$ gravity, due to nonvanishing $f_{R;i;j}$ (with $i \neq j$), the $\phi - \psi$ relation becomes scale dependent. Instead of the perturbation equation for the matter contrast δ , we provide here its evolution in term of the growth index $f = d \ln \delta / d \ln a$, that is the directly measured quantity at $z \sim 0.15$:

$$f'(a) - \frac{f(a)^2}{a} + \left[\frac{2}{a} + \frac{1}{a} E'(a) \right] f(a) - \frac{1 - 2Q}{2 - 3Q} \cdot \frac{3\Omega_m a^{-4}}{n E(a)^2 \tilde{R}^{n-1}} = 0, \quad (9)$$

$E(a) = H(a)/H_0$, \tilde{R} is the dimensionless Ricci scalar, and

$$Q = - \frac{2f_{RR} c^2 k^2}{f_R a^2}. \quad (10)$$

For $n = 1$ the previous expression gives the ordinary growth index relation for the Cosmological Standard Model. It is clear, from Eq.(9), that such a model suggests




Figure 2: Scale factor evolution of the growth index f : (*left*) modified gravity, in the case $\Omega_m = \Omega_{bar} \sim 0.04$, for the SNeIa best fit model with $n = 3.46$, (*right*) the same evolution in the case of a Λ CDM model. In the case of R^n -gravity it is shown also the dependence on the scale k . The three cases $k = 0.01, 0.001, 0.0002$ have been checked. Only the latter case shows a very small deviation from the leading behavior.

a scale dependence of the growth index which is contained into the corrective term Q so that, when $Q \rightarrow 0$, this dependence can be reasonably neglected. In the most general case, one can resort to the limit $aH < k < 10^{-3}h \text{ Mpc}^{-1}$, where Eq.(9) is a good approximation, and non-linear effects on the matter power spectrum can be neglected. Studying numerically Eq.(9), one obtains the growth index evolution in term of the scale factor; for the sake of simplicity, we assume the initial condition $f(a_{ls}) = 1$ at the last scattering surface as in the case of matter-like domination. The results are summarized in Fig.(2), where we show, in parallel, the growth index evolution in R^n -gravity and in the Λ CDM model. In the case of $\Omega_m = \Omega_{bar} \sim 0.04$, one can observe a strong disagreement between the expected rate of the growth index and the behavior induced by power law fourth order gravity models. These results seem to suggest that an extended gravity model which considers a simple power law of Ricci scalar, although cosmologically relevant at late times, is not viable to describe the evolution of Universe at all scales. In other words, such a scheme seems too simple to give account for the whole cosmological phenomenology. In fact, in Zhang 2006, a gravity Lagrangian considering an exponential correction to the Ricci scalar $f(R) = R + A \exp(-BR)$ (with A, B two constants), gives more interesting results and displays a grow factor rate which is in agreement with the observational results at least in the Dark Matter case. To corroborate this point of view, one has to consider that when the choice of $f(R)$ is performed starting from observational data (pursuing an inverse approach) as in Capozziello et al. 2005, the reconstructed Lagrangian is a non-trivial polynomial in term of the Ricci scalar. A result which directly suggests that the whole cosmological phenomenology can be accounted only with a suitable non-trivial function of the Ricci scalar rather than a simple power law function. As matter of fact, the results obtained with respect to the study of the matter power spectra in the case of R^n -gravity do not invalidate the whole approach, since they can be referred to the too simple form of the model.

3. DARK MATTER AS A CURVATURE EFFECT

The results obtained at cosmological scales motivates further analysis of $f(R)$ theories. In a sense, one is wondering whether the curvature fluid, which works as DE, can also play the role of effective DM thus yielding the possibility of recovering the observed astrophysical phenomenology by the only visible matter. It is well known that, in the low energy limit, higher order gravity implies a modified gravitational potential. Therefore, in our discussion, a fundamental role is played by the new gravitational potential descending from the given fourth order gravity theories we are referring to. By considering the case of a pointlike mass m and solving the vacuum field equations for a Schwarzschild-like metric, one gets from a theory $f(R) = f_0 R^n$, the modified gravitational potential Capozziello et al. 2007:

$$\Phi(r) = -\frac{Gm}{2r} \left[1 + \left(\frac{r}{r_c} \right)^\beta \right] \quad (11)$$

where

$$\beta = \frac{12n^2 - 7n - 1 - \sqrt{36n^4 + 12n^3 - 83n^2 + 50n + 1}}{6n^2 - 4n + 2} \quad (12)$$

which corrects the ordinary Newtonian potential by a power-law term. In particular, this correction sets in on scales larger than r_c which value depends essentially on the mass of the system. The corrected potential (11) reduces to the standard $\Phi \propto 1/r$ for $n = 1$ as it can be seen from the relation (12). The result (11) deserves some comments. As discussed in detail in Capozziello et al. 2007, we have assumed the spherically symmetric metric and imposed it into the field equations (1) considered in the weak field limit approximation. As a result, we obtain a corrected Newtonian potential which accounts for the strong non-linearity of gravity related to the higher-order theory. However, we have to notice that Birkhoff's theorem does not hold, in general, for $f(R)$ gravity but other spherically symmetric solutions than the Schwarzschild one can be found in these extended theories of gravity. The generalization of Eq.(11) to extended systems is achieved by dividing the system in infinitesimal mass elements and summing up the potentials generated by each single element. In the continuum limit, we replace the sum with an integral over the mass density of system taking care of eventual symmetries of the mass distribution (see Capozziello et al. 2007 for details). Once the gravitational potential has been computed, one may evaluate the rotation curve $v_c^2(r)$ and compare it with the data. For extended systems, one has typically to resort to numerical techniques, but the main effect may be illustrated by the rotation curve for the pointlike case, that is:

$$v_c^2(r) = \frac{Gm}{2r} \left[1 + (1 - \beta) \left(\frac{r}{r_c} \right)^\beta \right]. \quad (13)$$

Compared with the Newtonian result $v_c^2 = Gm/r$, the corrected rotation curve is modified by the addition of the second term in the r.h.s. of Eq.(13). For $0 < \beta < 1$, the corrected rotation curve is higher than the Newtonian one. Since measurements of spiral galaxies rotation curves signal a circular velocity higher than those which are predicted on the basis of the observed luminous mass and the Newtonian potential, the above result suggests the possibility that our modified gravitational potential may

fill the gap between theory and observations without the need of additional DM. It is worth noting that the corrected rotation curve is asymptotically vanishing as in the Newtonian case, while it is usually claimed that observed rotation curves are flat (i.e., asymptotically constant). Actually, observations do not probe v_c up to infinity, but only show that the rotation curve is flat within the measurement uncertainties up to the last measured point. This fact by no way excludes the possibility that v_c goes to zero at infinity. In order to observationally check the above result, we have considered a sample of LSB galaxies with well measured HI + H α rotation curves extending far beyond the visible edge of the system. LSB galaxies are known to be ideal candidates to test Dark Matter models since, because of their high gas content, the rotation curves can be well measured and corrected for possible systematic errors by comparing 21 - cm HI line emission with optical H α and [NII] data. Moreover, they are supposed to be Dark Matter dominated so that fitting their rotation curves without this elusive component is a strong evidence in favor of any successful alternative theory of gravity. Our sample contains 15 LSB galaxies with data on both the rotation curve, the surface mass density of the gas component and R -band disk photometry extracted from a larger sample selected by de Blok & Bosma 2002 . We assume the stars are distributed in an infinitely thin and circularly symmetric disk with surface density $\Sigma(r) = \Upsilon_\star I_0 \exp(-r/r_d)$ where the central surface luminosity I_0 and the disk scalelength r_d are obtained from fitting to the stellar photometry. The gas surface density has been obtained by interpolating the data over the range probed by HI measurements and extrapolated outside this range. When fitting to the theoretical rotation curve, there are three quantities to be determined, namely the stellar mass-to-light (M/L) ratio, Υ_\star and the theory parameters (β, r_c). It is worth stressing that, while fit results for different galaxies should give the same β , r_c is related to one of the integration constants of the field equations. As such, it is not a universal quantity and its value must be set on a galaxy-by-galaxy basis. However, it is expected that galaxies having similar properties in terms of mass distribution have similar values of r_c so that the scatter in r_c must reflect somewhat the scatter in the circular velocities. In order to match the model with the data, we perform a likelihood analysis determining for each galaxy, using, as fitting parameters β , $\log r_c$ (with r_c in kpc) and the gas mass fraction¹ f_g . As it is evident considering the results from the different fits, the experimental data are successfully fitted by the model (see Capozziello et al. 2007 for details). In particular, for the best fit range of β ($\beta = 0.80 \pm 0.08$), one can conclude that R^n gravity with $2.3 < n < 5.3$ (best fit value $n = 3.2$ which well overlaps the above mentioned range of n fitting SNeIa Hubble diagram) can be a good candidate to solve the missing matter problem in LSB galaxies without any Dark Matter. At this point, it is worth wondering whether a link may be found between R^n gravity and the standard approach based on Dark Matter haloes since both theories fit equally well the same data. As a matter of fact, it is possible to define an *effective Dark Matter halo* by imposing that its rotation curve equals the correction term to the Newtonian curve induced by R^n gravity. Mathematically, one can split the total rotation curve derived from R^n gravity as $v_c^2(r) = v_{c,N}^2(r) + v_{c,corr}^2(r)$ where the second term is the correction. Considering, for simplicity a spherical halo embedding a thin exponential disk, we may also write the total rotation curve as $v_c^2(r) = v_{c,disk}^2(r) + v_{c,DM}^2(r)$ with

¹This is related to the M/L ratio as $\Upsilon_\star = [(1 - f_g)M_g]/(f_g L_d)$ with $M_g = 1.4M_{HI}$ the gas (HI + He) mass, $M_d = \Upsilon_\star L_d$ and $L_d = 2\pi I_0 r_d^2$ the disk total mass and luminosity.




Figure 3: Best fit theoretical rotation curve superimposed to the data for the LSB galaxy NGC 4455 (left) and NGC 5023 (right). To better show the effect of the correction to the Newtonian gravitational potential, we report the total rotation curve $v_c(r)$ (solid line), the Newtonian one (short dashed) and the corrected term (long dashed).

$v_{c,disk}^2(r)$ the Newtonian disk rotation curve and $v_{c,DM}^2(r) = GM_{DM}(r)/r$ the Dark Matter one, $M_{DM}(r)$ being its mass distribution. Equating the two expressions, we get:

$$M_{DM}(\eta) = M_{vir} \left(\frac{\eta}{\eta_{vir}} \right) \frac{2^{\beta-5} \eta_c^{-\beta} (1-\beta) \eta^{\frac{\beta-5}{2}} \mathcal{I}_0(\eta) - \mathcal{V}_d(\eta)}{2^{\beta-5} \eta_c^{-\beta} (1-\beta) \eta^{\frac{\beta-5}{2}} \mathcal{I}_0(\eta_{vir}) - \mathcal{V}_d(\eta_{vir})}. \quad (14)$$

with $\eta = r/r_d$, $\Sigma_0 = \Upsilon_* i_0$, $\mathcal{V}_d(\eta) = I_0(\eta/2)K_0(\eta/2) \times I_1(\eta/2)K_1(\eta/2)^2$ and:

$$\mathcal{I}_0(\eta, \beta) = \int_0^\infty \mathcal{F}_0(\eta, \eta', \beta) k^{3-\beta} \eta'^{\frac{\beta-1}{2}} e^{-\eta'} d\eta' \quad (15)$$

with \mathcal{F}_0 only depending on the geometry of the system and “vir” indicating virial quantities. Eq.(14) defines the mass profile of an effective spherically symmetric Dark Matter halo whose ordinary rotation curve provides the part of the corrected disk rotation curve due to the addition of the curvature corrective term to the gravitational potential. It is evident that, from an observational viewpoint, there is no way to discriminate between this dark halo model and R^n gravity. Having assumed spherical symmetry for the mass distribution, it is immediate to compute the mass density for the effective dark halo as $\rho_{DM}(r) = (1/4\pi r^2)dM_{DM}/dr$. The most interesting features of the density profile are its asymptotic behaviors that may be quantified by the logarithmic slope $\alpha_{DM} = d \ln \rho_{DM}/d \ln r$ which can be computed only numerically as function of η for fixed values of β (or n). As expected, α_{DM} depends explicitly on β , while (r_c, Σ_0, r_d) enter indirectly through η_{vir} . The asymptotic values at the center and at infinity denoted as α_0 and α_∞ result particularly interesting. It turns out that α_0 almost vanishes so that in the innermost regions the density is approximately constant. Indeed, $\alpha_0 = 0$ is the value corresponding to models having an inner core such as the cored isothermal sphere and the Burkert model (Burkert 1995). Moreover, it is well known that galactic rotation curves are typically best fitted by cored dark halo models. On the other hand, the outer asymptotic slope is between -3 and -2 ,

²Here I_l and K_l , with $l = 1, 2$ are the Bessel functions of first and second type.

that are values typical of most dark halo models in the literature. In particular, for $\beta = 0.80$ one finds $(\alpha_0, \alpha_\infty) = (-0.002, -2.41)$, which are quite similar to the value for the Burkert model $(0, -3)$. It is worth noting that the Burkert model has been empirically proposed to provide a good fit to the LSB and dwarf galaxies rotation curves. The values of $(\alpha_0, \alpha_\infty)$ we find for the best fit effective dark halo therefore suggest a possible theoretical motivation for the Burkert-like models. Due to the construction, the properties of the effective Dark Matter halo are closely related to the disk one. As such, we do expect some correlation between the dark halo and the disk parameters. To this aim, exploiting the relation between the virial mass and the disk parameters, one can obtain a relation for the Newtonian virial velocity $V_{vir} = GM_{vir}/r_{vir}$:

$$M_d \propto \frac{(3/4\pi\delta_{th}\Omega_m\rho_{crit})^{\frac{1-\beta}{4}} r_d^{\frac{1+\beta}{2}} \eta_c^\beta V_{vir}^{\frac{5-\beta}{2}}}{2^{\beta-6}(1-\beta)G^{\frac{5-\beta}{4}} \mathcal{I}_0(V_{vir}, \beta)}. \quad (16)$$

We have numerically checked that Eq.(16) may be well approximated as $M_d \propto V_{vir}^a$ which has the same formal structure as the baryonic Tully-Fisher (BTF) relation $M_b \propto V_{flat}^a$ with M_b the total (gas + stars) baryonic mass and V_{flat} the circular velocity on the flat part of the observed rotation curve. In order to test whether the BTF can be explained thanks to the effective Dark Matter halo we are proposing, we should look for a relation between V_{vir} and V_{flat} . This is not analytically possible since the estimate of V_{flat} depends on the peculiarities of the observed rotation curve such as how far it extends and the uncertainties on the outermost points. For given values of the disk parameters, we therefore simulate theoretical rotation curves for some values of r_c and measure V_{flat} finally choosing the fiducial value for r_c that gives a value of V_{flat} as similar as possible to the measured one. Inserting the relation thus found between V_{flat} and V_{vir} into Eq.(16) and averaging over different simulations, we finally get:

$$\log M_b = (2.88 \pm 0.04) \log V_{flat} + (4.14 \pm 0.09) \quad (17)$$

while a direct fit to the observed data gives McGaugh 2005:

$$\log M_b = (2.98 \pm 0.29) \log V_{flat} + (3.37 \pm 0.13). \quad (18)$$

The slopes of the predicted and observed BTF are in good agreement thus leading further support to our approach. The zeropoint is markedly different with the predicted one being significantly larger than the observed one. However, it is worth stressing that both relations fit the data with similar scatter. A discrepancy in the zeropoint can be due to our approximate treatment of the effective halo which does not take into account the gas component. Neglecting this term, we should increase the effective halo mass and hence V_{vir} which affects the relation with V_{flat} leading to a higher than observed zeropoint. Indeed, the larger M_g/M_d is, the more the points deviate from our predicted BTF thus confirming our hypothesis. Given this caveat, we can conclude, with confidence, that R^n gravity offers a theoretical foundation even for the empirically found BTF relation. Although the results outlined along this paper are referred to a simple choice of fourth order gravity models ($f(R) = f_0 R^n$), they could represent an interesting paradigm. In fact, even if such a model is not suitable to provide the correct form of the matter power spectrum, as suggested by Troisi, A.: 2005, *Phys. Rev. D*, **71**, 043503. This suggests that a more

complicated Lagrangian is needed to reproduce the whole dark sector phenomenology at all scales, we have shown that considering extensions of GR can allow to explain some important issues of cosmological and astrophysical phenomenology. We have seen that extended gravity models can reproduce SNeIa Hubble diagram without Dark Matter, giving significant predictions even with regard to the age of Universe. In addition, the modification of the gravitational potential which arises as a natural effect in the framework of higher order gravity can represent a fundamental tool to interpret the flatness of rotation curves of LSB galaxies. Furthermore, if one considers the model parameters settled by the fit over the observational data on LSB rotation curves, it is possible to construct a phenomenological analogue of Dark Matter halo whose shape is similar to the one of the Burkert model. Since the Burkert model has been empirically introduced to give account of the Dark Matter distribution in the case of LSB and dwarf galaxies, this result could represent an interesting achievement since it gives a theoretical foundation to such a model. By investigating the relation among dark halo and the disk parameters, we have deduced a relation between M_d and V_{flat} which reproduces the baryonic Tully - Fisher. In fact, exploiting the relation between the virial mass and the disk parameters, one can obtain a relation for the virial velocity which can be satisfactorily approximated as $M_d \propto V_{vir}^a$. Even such a result seems very intriguing since it gives again a theoretical interpretation for a phenomenological relation. As a matter of fact, although not definitive, these results on $f(R)$ can represent a viable approach for future investigations and in particular support the quest for a unified view of the Dark Side of the Universe that could be interpreted as gravitational effects indeed.

References

- Burkert, A.: 1995, *Astroph. Journ.*, **447**, L25.
 Capozziello, S. et al.: 2003, *Int. J. Mod. Phys. D*, **12**, 1969.
 Capozziello, S., Troisi, A.: 2005, *Phys. Rev. D*, **72**, 044022.
 Capozziello S., Cardone V. F., Troisi A.: 2005, *Phys. Rev. D*, **71**, 043503.
 Capozziello, S., Cardone, V. F., Troisi, A.: 2007, *Mon. Not. Roy. Astron. Soc.*, **375**, 1423.
 Capozziello, S., De Filippis, E., Salzano, V.: 2009, *Mon. Not. Roy. Astron. Soc.*, **394**, 947.
 Capozziello, S., De Laurentis, M.: 2011, *Phys. Rept.*, **509**, 167.
 Copeland, E. J., Sami, M., Tsujikawa, S.: 2006, *Int. J. Mod. Phys. D*, **15**, 1753.
 Daly, R. A., Djorgovsky, S. G.: 2004, *Astrophys. J.*, **612**, 652.
 de Blok, W. J. G., Bosma, A.: 2002, *Astron. Astroph.*, **385**, 816.
 McGaugh, S. S.: 2005, *Astroph. Journ.*, **632**, 859.
 Zhang, P.: 2006, *Phys. Rev. D*, **73**, 123504.