

INVESTIGATION OF LS COUPLING IN OXYGEN III

B. Blagojević, M. V. Popović, Z. Pavlović and N. Konjević
Institute of Physics, 11080 Belgrade, P.O. Box 68 Yugoslavia

INTRODUCTION

LS or Russell-Saunders coupling is dominant for many transitions in the spectra of light elements. The spin-orbit interaction in atomic Hamiltonian becomes more important in comparison to the electrostatic separation between levels of the same principal quantum number n . Electrostatic separation increases as Z while the spin-orbit interaction grows as $Z^4\alpha^2$ where α is fine structure constant, so the LS-coupling scheme becomes inappropriate at some point. Systematic failure of the LS-coupling approximation is expected from lower to higher elements of an isoelectronic sequence for $nl-nl'$ transitions. The aim of this paper is to test accordance of experimental and theoretical data for LS-coupling in O III ions (3s-3p and 3p-3d transitions).

THEORY

Theoretical values calculated from multiconfiguration Dirac-Fock (MCDF) wave function of moderate accuracy are compared with our measured intensity ratios. For the case of pure LS coupling the relative line strength for a transition between levels J and J' is proportional to the factor (Cowan, 1981, Appendix I)

$$D^2 = (2J_1 + 1)(2J_2 + 1) \begin{Bmatrix} L_1 & S_1 & J_1 \\ J_2 & 1 & L_2 \end{Bmatrix} \quad (1)$$

Values of the $6j$ symbol are given in Appendix D of Cowan, 1981. The intensity ratio of two multiplet components is represented by (Glenzer et al., 1994)

$$\frac{I}{I'} = \frac{\lambda'^4 D^2_{J, J'} \frac{E' - E}{kT}}{\lambda^4 D'^2_{J, J'}} \quad (2)$$

where I , λ and I' , λ' are the total intensities and wavelengths of the two components, and E and E' are the energies of the upper levels of the two components, respectively.

EXPERIMENT

The light source was a low pressure pulsed arc with quartz discharge tube 10 mm internal diameter. The distance between aluminum electrodes was 16.2 cm and 3 mm diameter holes were located at the center of both electrodes to allow end-on plasma observations to be made. The central part around the pulsed arc axis was imaged 1 : 1 onto the entrance slit of the 1-m monochromator by means of the concave 1 m focal length, focusing mirror. A 30 mm diaphragm placed in front of the focusing mirror ensures that light comes from the narrow cone about the arc axis.

The monochromator with inverse linear dispersion 0.833 nm/mm in the first order of the diffraction grating, was equipped with the photomultiplier tube (PMT) and

a stepping motor. Signals from the PMT were led to a digital storage oscilloscope which was triggered by the voltage pulse from the Rogowski coil induced by the current pulse through the discharge tube. The discharge was driven by a 15.2 μF low inductance capacitor charged to 3 KV (peak current 15 KA) and fired by an ignitron. The stepping motor and oscilloscope are controlled by personal computer which was also used for data acquisition. Recordings of spectral line shapes were performed shot-by-shot. At each wavelength position of the monochromator time evolution and decay of the plasma radiation were recorded and memorized by the oscilloscope. Eight such signals were averaged at each wavelength. Further, to construct the line profiles these averaged signals at different wavelengths and at various times of the plasma existence were used to construct line profiles. Greatest care was taken to find the optimum conditions with the least line self absorption. It was found that the percentage of oxygen in the mixture was of crucial importance for the elimination of self-absorption. The ratio $\text{O}_2 : \text{He} = 0.6 : 99.4$ was determined after a number of experiments in which O_2 was diluted gradually until strong line intensities O III are found proportional to the concentration of O_2 in the gas mixture. During the spectral line recording continuous flow of oxygen-helium mixture was maintained at a pressure of about 400 [Pa].

PLASMA DIAGNOSTICS

For the electron density measurements we use the width of He II P_α 468.6 nm line. The full width at half maximum $\Delta\lambda_{\text{FWHM}}$ of this line is related to the electron density N_e using the following relationship (Pittman and Fleurier, 1982; 1986; Fleurier and Le Gall, 1984)

EXPERIMENTAL RESULTS AND DISCUSSION

The experimental results for intensity ratios R_m from $3s\text{-}(^2P^o)3p$ multiplet ($I_{2,3}/I_{1,2}$) and $3p\text{-}(^2P^o)3d$ multiplet ($I_{3,4}/I_{2,3}$) of O III ions are compared with theoretical ratios R_{th} given in Table 1 together with electron concentration. Theoretical ratios are $R_{th} = 1.84$ for $3s\text{-}(^2P^o)3p$ multiplet and $R_{th} = 1.43$ for $3p\text{-}(^2P^o)3d$ multiplet. Table 1 shows that LS coupling approximation is fulfilled for investigated transitions in O III ions.

Electron density N_e [10^{17} cm $^{-3}$]	Transition array	
	$3s\text{-}(^2P^o)3p$ R_m / R_{th}	$3p\text{-}(^2P^o)3d$ R_m / R_{th}
0.64	1.07	1.13
1.05	1.14	1.06
1.22	0.96	1.14
1.32	1.00	1.05
1.33	0.95	0.94
1.30	1.03	1.01
1.21	0.95	0.97
1.07	0.96	0.95
0.90	0.94	0.95
0.73	0.94	0.90
0.57	0.98	1.10
0.46	0.97	1.01
0.37	0.84	1.10
	<0.98>	<1.02>

Table 1. R_m / R_{th} ratios for $3s\text{-}(^2P^o)3p$ and $3p\text{-}(^2P^o)3d$ transitions in O III. Averaged ratios for measured electron densities N_e are given in <> brackets

REFERENCES

- Cowan R.D., 1981, The Theory of Atomic Structure and Spectra, (University of California Press, Berkeley).
- Fleurier C. and Le Gall P., 1984, J.Phys.B 17, 4311.
- Glenzer S., Kunze H.-J., Musielok J., Kim Y.-K. and Wiese W.L., 1994, Phys.Rev.A 49, 221.
- Griem H.R. and Shen K.Y., 1961, Phys.Rev. 122, 1490.
- Kobilarov R., Konjević N., and Popović M.V., 1981, Phys.Rev.A 40, 3871.
- Konjević N. and Wiese W.L., 1976, J.Phys.Chem.Ref.Data 5, 259.
- Pittman T.L. and Fleurier C., 1982, in Proceedings of the Sixth International Conference on Spectral Line Shapes, 4 Boulder, (de Gruyter, Berlin, 1983).
- Pittman T.L. and Fleurier C., 1986, Phys.Rev A 33, 1291.
- Wiese W.L., 1965, in Plasma Diagnostic Techniques, edited by Huddleston R.M. and Leonard S.L. (Academic, New York).
- Wiese W.L., Fuhr J.R. and Deters T.M., 1996, J.Phys.Chem.Ref.Data, Monography No.7.